

Extended summaries

Neurotox '98

The following are Extended Summaries based on posters presented at the meeting 'Neurotox '98' organised by Df Beadle, IS Blagborough, MG Ford, R Greenwood, Df Ray and PN Usherwood on behalf of the Physicochemical and Biophysical Panel of the SCI Pesticides Group and held at St Catherine's College, Oxford, on 28–31 July 1998. They are entirely the responsibility of the authors and do not necessarily reflect the views of the Editorial Board of Pesticide Science.

Can selective peptides be combined efficiently with agrochemicals? A new approach to insect control

Dalia Gordon*

CEA, Département d'Ingénierie et d'Etudes des Protéines, Bat 152, C E Saclay, F- 91191 Gif sur Yvette, France

Abstract: Sodium channels have been a major target for the development of insecticides such as synthetic pyrethroids. However, insecticides currently available induce resistance and present limited selectivity to insect pests. Molecular and biochemical studies, as well as binding experiments using radiolabelled neurotoxins, have shown that sodium channels expressed in various insect orders must be structurally and pharmacologically different. At least three groups of peptide neurotoxins derived from scorpion venom are highly active on insects and very weakly or practically inactive on mammals. It is proposed that various insecticides are examined for possible cooperative interactions with the peptide toxins highly active on insects, and pairs of ligands are identified that will increase the selectivity not only between mammals and insects but also between different pest and non-pest insects. This is feasible on the basis of the differential allosteric modulations observed between Lqh α IT, an α -toxin highly active on insects, and brevetoxin on locust *versus* cockroach and rat brain sodium channels. Moreover, combination of Lqh α IT with the pyrethroid deltamethrin increased the binding of [¹²⁵I] Lqh α IT by more than 1.8-fold, and the combined presence of brevetoxin further increased the binding. Such allosteric modulation may provide a new approach to increase the selective activity of pesticides on target organisms by simultaneous application of allosterically interacting drugs, designed on the basis of the selective peptide toxins.

Keywords: insect sodium channel; scorpion toxin; allosteric interactions; neurotoxin receptor site; insect control; pyrethroids

INTRODUCTION

The voltage-gated sodium channels are critical elements in nerve excitability and serve as the initial target for many toxins developed by a wide variety of venomous animals, including scorpions, spiders and sea anemones. Sodium channels have been a major target for the development of insecticides, such as synthetic pyrethroids.¹ However, insecticides currently available present limited selectivity to insect pests, induce resistance in the pest insects and may disturb the delicate balance between natural predators and pests. Thus, a search for new targets and highly selective insecticides is an important goal in insect control management. Information on the variability in the structure of insect sodium channels and ligands which are highly active on insects may offer a new approach for the strategy of insect control.

Diversity of sodium channels in various insects

We have previously shown, using selective immunoprecipitation of insect sodium channels by anti-peptide antibodies, that sodium channel α subunit proteins from various insect neuronal membranes are similar in their biochemical properties to rat brain sodium channels. Sodium channels of locust, cockroach, blowfly head and lepidopteran larvae revealed apparent molecular masses of 245–260 kDa, which were devoid of a disulfide-linked smaller subunit, in contrast to rat brain sodium channels.^{2,3} Cloning of *Drosophila* sodium channel,⁴ partial proteolysis of various insect sodium channel proteins³ as well as binding experiments using radiolabelled neurotoxins^{5–7} have shown that sodium channels expressed in various insect orders must be structurally and pharmacologically different. Binding studies using [¹²⁵I] Lqh α IT (Lqh – *Leiurus quinquestriatus hebraeus*; IT – insect toxin), the scorpion α -toxin most active on insects, and competition binding studies with several scorpion toxins and the sea anemone toxin ATX II demonstrated that the affinities of the different toxins to locust and cockroach neuronal membranes are different, suggesting structural differences in the sodium channel receptor sites.⁶ Dissociation rate constants of bound [¹²⁵I] AahIT (Aah (called also AaH) – *Androctonus australis* Hector), the excitatory insect selective toxin, in several different insect neuronal membranes revealed k_{off} values in the range

* Correspondence to: Dalia Gordon, CEA, Département d'Ingénierie et d'Etudes des Protéines, Bat 152, C E Saclay, F-91191 Gif sur Yvette, France
(Received 10 August 1998; revised version received 25 November 1998; accepted 20 May 1999)




Figure 1. Allosteric interactions among several neurotoxin receptor sites on sodium channels. Interactions between the lipid soluble polyether toxin, brevetoxin PbTx-1, and scorpion α -toxins on rat brain, cockroach (*Periplaneta americana*) and locust (*Locusta migratoria*) sodium channels (Top). The lower panel presents a cartoon of a section through sodium channel α -subunit to demonstrate several allosteric interactions observed by binding studies on the locust sodium channel. The ovals represent the distinct neurotoxin receptor sites as indicated. The arrows indicate the allosteric modulation by binding of one ligand to its receptor site on the binding of the another ligand, linked by the arrows. (+) sign indicates a positive co-operative effect. From References 6 and 21.

of 1.33×10^{-3} (in cockroach) to $98.8 \times 10^{-3} \text{ s}^{-1}$ (in *Spodoptera* larvae), further indicating that the different insect sodium channels must be structurally distinct.⁷ These differences provide the possibility of targetting different sodium channels by selective ligands.

Diversity in sodium channel ligands

At least three groups of peptide neurotoxins derived from scorpion venom are highly active on insects and very weakly or practically inactive on mammals.^{8–10,23} The scorpion α -toxin Lqh α IT is highly active on insects,^{6,10} and the two anti-insect selective toxins, the excitatory (AahIT) and depressant (LqhIT2),^{8,11} bind to distinct, high-affinity receptor sites on insect sodium channels and modify their activity differently.^{5,6,10,12–16} The high activity and selectivity of the peptide toxins to insects provide the possibility to use them as leads to develop highly selective insecticides. However, a detailed functional analysis of these toxins and an understanding of the structural basis of

their selectivity to insects must be obtained as a prerequisite for modifying the peptides in a rational manner. These studies have started with the recent functional expression and analysis by mutagenesis of Lqh α IT^{16–19} and the depressant and excitatory toxins by Gurevitz and co-workers (Ref 22 and unpublished).

Combinations of ligands

The proposed approach takes advantage of the availability of many ligands that specifically target voltage-gated sodium channels in general, and insect sodium channels in particular. At least seven distinct classes of neurotoxins have been designated, on the basis of physiological activity and competitive binding studies, that bind to several distinct receptor sites on the sodium channel protein.²⁰ Although the sodium channel receptor sites are topologically separated, there are strong allosteric interactions among them. We have shown that the lipid-soluble sodium channel activators, veratridine and brevetoxin, reveal divergent allosteric modulation of the binding of scorpion α -toxins at homologous receptor sites on mammalian and insect sodium channels.^{21,23}

Brevetoxin reveals negative allosteric interaction with Aah II on rat brain sodium channels, but increases the affinity of Lqh α IT to locust sodium channels, and has no significant effect on Lqh α IT binding in cockroach neuronal membranes (Fig 1). The differences suggest functionally important distinctions between these channel subtypes.

In this approach it is proposed to examine various insecticides, including pyrethroids, for possible co-operative interactions with the peptide toxins highly active on insects, and to identify pairs (or more) of ligands that will increase the selectivity not only between mammals and insects but also between different pest and non-pest insects. This is feasible on the basis of the differential allosteric modulations observed above. To exemplify this approach, the scorpion α -toxin Lqh α IT was combined with delta-methrin. As expected, the binding of [¹²⁵I]Lqh α IT was increased by more than 1.8-fold by the pyrethroid, and the combined presence of veratridine and/or brevetoxin further increased the binding (Gilles and Gordon, unpublished). The allosteric modulation may provide a new approach to increase selective activity of pesticides on target organisms by simultaneous application of allosterically interacting drugs, designed on the basis of the selective toxins.

The most recent advances in cloning, characterization and functional expression of new scorpion toxins highly active on different pest insects, the structural and mutagenesis studies that are currently in progress may provide a new momentum to the clarification of the structural basis of insect selectivity of the peptide toxins and enable rational drug design of highly selective insecticides.²² The combined approach, using pairs of the novel ligands and known agrochemicals, may enable significant reduction in the amount of insecticides used in practice and may aid in

preventing the development of resistance in pest insects.

REFERENCES

- Bloomquist JR, Ion channels as target for insecticides. *Ann Rev Entomol* 41:163–190 (1996).
- Gordon D, Merrick D, Wollner DA and Catterall WA, Biochemical properties of sodium channels in a wide range of excitable tissues studied with site-directed antibodies. *Biochemistry* 27:7032–7038 (1988).
- Gordon D, Moskowitz H and Zlotkin E, Sodium channel polypeptides in central nervous systems of various insects identified with site-directed antibodies. *Biochim Biophys Acta* 1026:80–86 (1990).
- Loughney K, Kreber R and Ganetzky B, Molecular analysis of the *para* locus, a sodium channel gene in *Drosophila*. *Cell* 58:1143–1154 (1989).
- Moskowitz H, Herrmann R, Zlotkin E and Gordon D, Variability among insect sodium channels revealed by binding of selective neurotoxins. *Insect Biochem Molec Biol* 24:13–19 (1994).
- Gordon D, Martin-Eauclaire MF, Cestèle S, Kopeyan C, Carlier E, Ben Khalifa R, Pelhate M and Rochat H, Scorpion toxins affecting sodium current inactivation bind to distinct homologous receptor sites on rat brain and insect sodium channels. *J Biol Chem* 271:8034–8045 (1996).
- Fishman L, Herrmann R, Gordon D and Zlotkin E, Insect tolerance to neurotoxic polypeptide: pharmacokinetic and pharmacodynamic aspects. *J Exp Biol* 200:1115–1123 (1997).
- Zlotkin E, Miranda F and Rochat H, Chemistry and pharmacology of Buthinae scorpion venoms, in *Arthropod Venoms*, ed. by Bettini S, Springer Verlag pp 317–369 (1978).
- Lester D, Lazarovici P, Pelhate M and Zlotkin E, Purification, characterization and action of two insect toxins from the venom of the scorpion *Buthotus judaicus*. *Biochim Biophys Acta* 701:370–381 (1982).
- Eitan M, Fowler E, Herrmann R, Duval A, Pelhate M and Zlotkin E, A scorpion venom neurotoxin paralytic to insects that affects sodium current inactivation: purification, primary structure, and mode of action. *Biochemistry* 29:5941–5947 (1990).
- Zlotkin E, Kadouri D, Gordon D, Pelhate M, Martin MF and Rochat H, An excitatory and a depressant insect selective toxin from scorpion venom both affect sodium conductance and possess a common binding site. *Arch Biochem Biophys* 240:877–887 (1985).
- Gordon D, Jover E, Couraud F and Zlotkin E, The binding of an insect selective neurotoxin (AaIT) from scorpion venom to locust synaptosomal membranes. *Biochim Biophys Acta* 778:349–358 (1984).
- Gordon D, Moskowitz H, Eitan M, Warner C, Catterall WA and Zlotkin E, Localization of receptor sites for insect-selective toxins on sodium channels by site-directed antibodies. *Biochemistry* 31:7622–7628 (1992).
- Gordon D and Zlotkin E, Binding of an alpha scorpion toxin to insect sodium channels is not dependent on membrane potential. *FEBS Lett* 315:125–129 (1993).
- Zlotkin E, Eitan M, Bindokas V, Adams ME, Moyer M, Brukhart W and Fowler E, Functional duality and structural uniqueness of depressant insect-selective neurotoxins. *Biochemistry* 30:4814–4820 (1991).
- Gordon D, Savarin P, Gurevitz M and Zinn-Justin S, Functional anatomy of scorpion toxins affecting sodium channels. *J Toxicol Toxin Rev* 17:131–158 (1998).
- Zilberberg N, Gordon D, Pelhate M, Adams ME, Norris F, Zlotkin E and Gurevitz M, Functional expression and genetic modification of an alpha scorpion neurotoxin. *Biochemistry* 35:10215–10222 (1996).
- Zilberberg N, Froy O, Loret E, Cestèle S, Arad D, Gordon D and Gurevitz M, Identification of structural elements of a scorpion α -neurotoxin important for receptor site recognition. *J Biol Chem* 272:14810–14816 (1997).
- Tugarinov V, Kustanovich I, Zilberberg N, Gurevitz M and Anglister J, Solution structure of a highly insecticidal recombinant scorpion α -toxin and a mutant with increased activity. *Biochemistry* 36:2414–2424 (1997).
- Gordon D, Sodium channels as targets for neurotoxins: mode of action and interaction of neurotoxins with receptor sites on sodium channels, in *Toxins And Signal Transduction*, ed by Lazarowici P and Gutman Y, Harwood Press, Amsterdam. pp 119–149 (1997).
- Cestèle S, Ben Khalifa R, Pelhate M, Rochat H and Gordon D, α -Scorpion toxins binding on rat brain and insect sodium channels reveal divergent allosteric modulations by brevetoxin and veratridine. *J Biol Chem* 270:15153–15161 (1995).
- Oren DA, Froy O, Amit E, Kleinberger-Doron N, Gurevitz M and Shaanan B, An excitatory scorpion toxin with a distinctive feature: an additional α helix at the C terminus and its implications for interaction with insect sodium channels. *Structure* 6:1095–1103 (1998).
- Gordon D, A new approach to insect-pest control – combination of neurotoxins interacting with voltage-sensitive sodium channels to increase selectivity and specificity. *Invertebr Neurosci* 3:103–116 (1997).

Characterisation of the relationship between binding sites for imidacloprid and other nicotinic ligands in insects

Robert J Lind,^{1*} Martin S Clough,² Fergus GP Earley,² Susan Wonnacott¹ and Stuart E Reynolds¹

¹Department of Biology and Biochemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK

²Zeneca Agrochemicals, Jealott's Hill Research Station, Bracknell, Berkshire, RG42 6ET, UK

Abstract: Radioligand binding studies using the neurotoxins α -bungarotoxin, epibatidine, imidacloprid (IMI) and methyllycaconitine reveal heterogeneity at the level of the nicotinic acetylcholine receptor (nAChR) in membranes from the peach potato aphid *Myzus persicae* (Sulzer) and further suggest the presence of more than one ligand binding site per nAChR. These sites are able to interact allosterically with each other. Of particular interest, [³H]IMI has over an order of magnitude higher affinity in membranes of hemipteran pest species than in non-hemipteran insects, which may help explain why IMI is particularly effective for the control of sucking pests.

Keywords: *Myzus persicae*; nicotinic acetylcholine receptor; imidacloprid; methyllycaconitine; epibatidine; radioligand binding

* Correspondence to: Robert J Lind, Zeneca Agrochemicals, Jealott's Hill Research Station, Bracknell, Berks RG42 6ET, UK
E-mail: rob.lind@aguk.zeneca.com

Contract/grant sponsor: Biotechnology and Biological Science Research Council

Contract/grant sponsor: Zeneca Agrochemicals

(Received 10 August 1998; accepted 20 May 1999)